

# Searching for exceptional gravitational-wave sources in the LIGO-Virgo-KAGRA (LVK) data

- Model-independent searches
  - Core-collapse supernovae
  - LVK Workshop in Warsaw

Marek Szczepańczyk

Department of Physics, University of Warsaw


PAiP-2025 conference  
Warsaw, 20.02.2025

# Return to Poland

- Ph.D., ~5 years: Embry-Riddle Aeronautical University (Arizona)
- Postdoc, ~5 years: University of Florida
- Assistant Professor, present: University of Warsaw (permanent position and a Polish Returns grant)

Homepage: <https://www.fuw.edu.pl/~mszczepancyk/>

Prof. Jerzy Lewandowski was my Inviting Scientist for Polish Returns grant




# Exceptional GW sources

**Exceptional astrophysical sources might play an important role in our endeavor of exploring the Universe.**


- **New GW source populations:**
  - Compact binaries: binaries with eccentric orbits, hyperbolic encounters, head-on collisions, extreme mass ratio, sub-solar mass binaries
  - GW bursts: core-collapse supernovae, neutron star or pulsar glitches, cosmic strings
- **Multi-messenger GW sources (electromagnetic waves, neutrinos, cosmic rays):** BNS, NSBH, BNS post-merger
- **GW sources with new phenomena (usually weaker effects):**
  - GR: pre- and post-merger higher harmonics, GW cross-polarization, black hole kicks, GW memory, effects of precession, high spins, black hole formation, lensed binaries
  - Beyond GR: GW echo, beyond-quadrupolar GW polarizations,

# Model-independent searches

- **Coherent WaveBurst** (cWB, Klimenko+16) is a software designed to detect a wide range of burst transients without prior knowledge of the signal morphology
- cWB uses minimal assumptions, for example growing frequency over time in case of binaries
- **Complementing template-based searches**
- cWB has detected:
  - **GW150914** - the very first GW (PRL 116, 061102)
  - **GW190521** - an intermediate mass binary black hole (PRL 125, 101102)
  - It regularly detects GWs together with template-based searches
- The cWB contributes results to several LVK papers during each observing run.




<https://gwburst.gitlab.io/>



# Model-independent searches classification


## Compact binary searches (minimally modeled)

Binary black holes  
Binary neutron stars  
Black hole - neutron star



e.g. Mishra+23 ([2201.01495](#))


Binaries with eccentric orbits  
Intermediate-mass black holes  
Hyperbolic encounters  
Extreme mass-ratio



e.g. MS+21 ([2009.11336](#))


## Generic searches (unmodeled)

Core-collapse supernovae  
Pulsar glitches  
Cosmic strings  
Unknown



e.g. MS+24 ([2305.16146](#))


## Low-latency searches



Public alerts for  
multi-messenger observations:  
electromagnetic, cosmic rays,  
and neutrino

e.g. Chaudhary+24 ([2308.04545](#))

## Searches for new phenomena



Higher harmonics  
GW cross-polarization  
Deviations from GR

e.g. Vedovato+22 ([2108.13384](#))


# Core-Collapse Supernova (CCSN)

*Nova on the sky!*



1-2 per century in Milky Way (?)

- Burning of a star:  $H \rightarrow He \rightarrow \dots \rightarrow Fe$
- After exceeding Chandrasekhar mass of  $1.4 M_{\odot}$  the iron core collapses.
- 99% of explosion energy escapes with neutrinos!

**Explosion mechanism(s)  
is still unknown**






Janka+12



SN 1054 (Crab Nebula)

# When will we discover GWs? (realistically: Galactic CCSN)



# LVK and CCSN Theory

Example: Mezzacappa et al 2023

- CCSNe are the most challenging astronomical events to model:
  - All four fundamental forces are important
  - Neutrino transport
  - Computational challenges
- Last joint workshop between LVK and CCSN modelers was at Caltech in 2017
  - Creating Supernova Multimessenger Consortium
- Agenda/webpage: work in progress

**LVK workshop:  
July 21-23, 2025, in Warsaw**

Note: it's right after the GR24/Amaldi16 meeting in Glasgow (July 14-18, 2025)





Image: Michael Sandoval, ORNL








# Summary

- Model-independent searches
  - Preparing for exceptional/special GW sources
- Core-Collapse Supernova (CCSN)
  - Next Galactic CCSN: one of the most interesting events of the century
- Joint workshop between LVK and CCSN modelers:  
July 21-23, 2025 in Warsaw

More info: <https://www.fuw.edu.pl/~mszczepanczyk/>

# Extras


# coherent WaveBurst (cWB)




# Wavescan

- Wavescan (Klimenko+22, [2201.01096](https://arxiv.org/abs/2201.01096)): high-resolution time-frequency transform
- Heisenberg rule for signal processing:  $\sigma_t^2 \sigma_\omega^2 \geq \frac{1}{4}$ 
  - Multiresolution analysis and wavelet stack
- Wavescan transform combines the maps from different resolution into a single time-frequency map
  - Spectral and temporal leakage is minimized.

## Wavelet stack






Klimenko, 2021  
Arxiv 2201.01096



# Eccentric binaries

- Eccentric binaries: compact binaries elliptical orbits.
  - Dynamical formation
- Bhaumik et al (MS) 2024 ([2410.15192](https://arxiv.org/abs/2410.15192))
  - Comparison between waveform models
  - Sensitivity studies and recommendations
- Mishra et al (MS) 2024 ([2410.15191](https://arxiv.org/abs/2410.15191))
  - O3 data reanalysis
  - 3 new GWs: consistent with stellar BHs, one event has large mass-ratio (possible dynamic formation)



# O4 cWB low-latency searches

- The cWB searches: cWB-AllSky (generic) and cWB-BBH
- Analysis:
  - LH: searches, significance
  - LHV: sky map follow-up

## cWB-AllSky (generic)



- cWB-XP and cWB-2G
- Public alert for GW bursts: “fluences” (~luminosity), peak frequency, duration
- Only one event so far [S200114f](#) (O3) - classified as noise offline

## cWB-BBH search

- cWB-BBH events are treated as CBC (RODA: [M2200164](#))
- 3 events so far
- **It’s capable to detect “vanilla” and special/exceptional compact binaries**
- Complementing matched filtering
- It detects around **80%** of BBHs identified by matched filtering searches (HL network)



# GW190521

- Intermediate-mass black holes (IMBHs) - between stellar mass ( $100 M_{\odot}$ ) and supermassive ( $10^5 M_{\odot}$ ). The origin is not yet well understood.
  - Probing pair-instability mass gap (Stars with He mass in ( $64 M_{\odot}$ ,  $135 M_{\odot}$ ))
  - Formation channels
  - Most distant GW sources
- GW190521 - first conclusive evidence of an IMBH.
- No chirping structure
- Detection significance (see MS+21, [2009.11336](#)):
  - Online: 1 per 28 years
  - Offline: 1 per 4900 years (established by cWB)
  - Challenges: scatter noise, blips



# Optically targeted searches

- While waiting for a Galactic CCSN, we can systematically constrain its engine with CCSNe at MPc range -> optically targeted searches
- O1-O2 search (Abbott+19, [1908.03584](#)):
  - First observational constraints of a CCSN engine (my main PhD thesis result)
- O3 search (Szczepanczyk+24, [2305.16146](#)):
  - We could not beat previous limits
- SN 2023ixf search (Abac+24, [2410.16565](#), special O4 paper):
  - GW energy emission: constraints improved by an order of magnitude






# Parameter Estimation

Recently a lot of efforts to extract physical parameters from CCSN. See review in Mezzacappa&Zanolin+24 ([2401.11635](#)), examples:

- Proto-neutron star (PNS) evolution: Casallas-Lagos+23 ([2304.11498](#)), Bizouard+21 ([2012.00846](#)),
- Equation of State: Edwards+21 ([2009.07367](#)),
- SN kicks (GW memory): Richardson+21 ([2109.01582](#))
- Standing Accretion Shock Instability: Takeda+21 ([2107.05213](#))
- PNS rotation: Chan+21 ([ADS](#)), Hayama+18 ([1802.03842](#))
- Rotation properties: Pastor-Marcos+23 ([2308.03456](#)), Villegas+23 ([2304.01267](#))

## Non rotating scenario



↑ Bounce time is determined by  $\nu$  observation