



Contribution ID: 21

Type: **Talk**

Dynamical models of the Milky Way nuclear star cluster

Monday, 19 August 2024 14:40 (20 minutes)

The nuclear star cluster (NSC) of the Milky Way has been extensively studied in the last decades, using ground-based astrometry and spectroscopy of ~ 10000 stars in the inner 10 pc.

The Galactic centre is unique in that we have a direct measurement of the mass of the supermassive black hole (SMBH) from the motion of S-stars, which can be tracked for a significant fraction of the orbital period. However, the remaining vast majority of stars in the NSC are essentially a single kinematic snapshot, and fall within the realm of classical methods for inferring the gravitational potential from the assumption of dynamical equilibrium.

I present a new twist in this analysis based on the iterative self-consistent modelling method with action-space distribution functions. I discuss the constraints on the NSC properties and the intrinsic degeneracies and limitations of this inference procedure, which are relevant more generally for dynamical modelling of extragalactic nuclear clusters.

Affiliation

University of Surrey

Current Position

Postdoc

Primary author: VASILIEV, Eugene (University of Surrey)

Presenter: VASILIEV, Eugene (University of Surrey)

Session Classification: Nuclear star clusters and the Galactic nucleus

Track Classification: Numerical approaches to modelling stellar systems and their constituents