



Contribution ID: 11

Type: **Talk**

Star cluster properties from intermediate-mass black hole mergers

Friday, 23 August 2024 10:00 (20 minutes)

Next-generation (XG) ground-based gravitational wave observatories are expected to be sensitive to mergers of intermediate-mass black holes (IMBHs) with a total source-frame mass in the hundreds out to a high redshift. Runaway tidal encounters lead to the formation of IMBHs in the cores of dense stellar clusters. I will discuss how single IMBH–IMBH merger events, occurring after the coalescence of clusters that sank into the center of their host galaxies, can be used to infer the properties of their progenitor star clusters. Implementing an astrophysically motivated analytic model and performing binary parameter estimation for our massive events in a network of three XG detectors, we find that inferring the structural properties of clusters through this channel is challenging due to model degeneracy. Nonetheless, the redshifts of cluster formation are better measured, and the cluster formation history may be inferred from an observed population of IMBH–IMBH mergers.

Affiliation

Johns Hopkins University

Current Position

PhD Student

Primary authors: Prof. BERTI, Emanuele (Johns Hopkins University); Prof. ANTONINI, Fabio (Cardiff University); Dr NG, Ken (Johns Hopkins University); KRITOS, Konstantinos (Johns Hopkins University); Mr REALI, Luca (Johns Hopkins University)

Presenter: KRITOS, Konstantinos (Johns Hopkins University)

Session Classification: Chris Belczynski memorial session on compact objects and gravitational wave sources

Track Classification: Chris Belczynski memorial session on compact objects and gravitational wave source