



Contribution ID: 102

Type: **Talk**

## Formation of globular clusters with intermediate-mass black holes

*Tuesday, 20 August 2024 17:34 (3 minutes)*

The formation process of star clusters has not been fully understood yet. Recent numerical simulations of star-cluster formation are reaching the mass of globular clusters ( $1e6$  Msun). However, in hydrodynamics simulation with N-body, the stars have been treated as super particles, representing several stars as one particle. We have developed a new N-body/smoothed-particle hydrodynamics (SPH) code, ASURA+BRIDGE, and have performed N-body/SPH simulations of forming star clusters with almost  $1e6$  Msun by resolving individual stars. Our code can integrate the orbits of stars without gravitational softening using a direct-tree hybrid N-body code, PeTar (Wang et al. 2020), combined with our N-body/SPH simulation code, ASURA+BRIGE (Fujii et al. 2021). In our simulations, runaway collisions of stars occurred in the forming globular clusters, and the mass of the very massive stars (VMSs) formed via runaway collisions reached a maximum of  $1e4$  Msun. According to a stellar evolution model, they can collapse to IMBHs with a mass of a few thousand Msun. Our results suggest that some globular clusters may host an IMBH more massive than 1000 Msun. In addition, such VMSs quickly lose their mass via stellar wind and pollute the surrounding gas. In our simulations, so-called second population stars are also formed from the polluted gas. They typically have more than 0.1% of the mass originating from the VMS.

### **Affiliation**

The University of Tokyo

### **Current Position**

Senior Scientist or Faculty

**Primary author:** FUJII, Michiko (The University of Tokyo)

**Presenter:** FUJII, Michiko (The University of Tokyo)

**Session Classification:** Virtual Poster Presentations